

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings of claims in the application:

Listing of Claims:

1. (Previously presented) A method of increasing the mutation rate of a retrovirus or flavivirus, comprising administering an RNA nucleoside analog to a virally infected cell, wherein the analog is incorporated by a polymerase into an RNA copy of a genomic nucleic acid encoding the retrovirus or flavivirus, said analog replacing a first natural occurring nucleotide having a first complementary nucleotide wherein said analog complements a second nucleotide which is other than the first nucleotide, thereby inducing the retrovirus or flavivirus to mutate.
2. (Original) The method of claim 1, wherein the RNA nucleoside analog replaces uracil.
3. (Original) The method of claim 1, wherein the RNA nucleoside analog replaces adenine.
4. (Original) The method of claim 1, wherein the RNA nucleoside analog replaces cytidine.
5. (Original) The method of claim 1, wherein the RNA nucleoside analog replaces guanine.
6. (Original) The method of claim 1, wherein the RNA nucleoside analog is incorporated by the polymerase into the RNA copy of the genomic nucleic acid with an efficiency at least about 0.1% that of a naturally occurring complementary nucleic acid.

7. (Currently amended) The method of claim 1, wherein the method further includes the proviso that the RNA nucleoside analog is not ribavirin or a 5-halo analog of ~~1- β -D-ribofuranosylimidazole-4-carboxamide~~ 1- β -D-ribofuranosylimidazole-4-carboxamide.

8. (Original) The method of claim 1, wherein the RNA analog is a non-chain terminating analog.

9. (Original) The method of claim 1, wherein the method further includes the proviso that if the virus is HIV, then the RNA nucleoside analog is not HEPT or a 2',5'-bis-O-sialylated-3'-spiro-substituted (TSAO) adenine, hypoxanthine, N¹-alkyl-hypoxanthine, or xanthine or a nucleoside analog that is incorporated and extended at high efficiency by reverse transcriptase of HIV.

10. (Original) The method of claim 1, wherein the nucleoside analog is selected from the group consisting of N⁴-aminocytidine, N¹-methyl-N⁴-aminocytidine, 3,N⁴-ethenocytidine, 3-methylcytidine, 5-hydroxycytidine, N⁴-dimethylcytidine, 5-(2-hydroxyethyl)cytidine, 5-chlorocytidine, 5-bromocytidine, N⁴-methyl-N⁴-aminocytidine, 5-aminocytidine, 5-nitrosocytidine, 5-(hydroxyalkyl)-cytidine, 5-(thioalkyl)-cytidine and cytidine glycol, 5-hydroxyuridine, 3-hydroxyethyluridine, 3-methyluridine, O²-methyluridine, O²-ethyluridine, 5-aminouridine, O⁴-methyluridine, O⁴-ethyluridine, O⁴-isobutyluridine, O⁴-alkyluridine, 5-nitrosouridine, 5-(hydroxyalkyl)-uridine, and 5-(thioalkyl)-uridine, 1,N⁶-ethenoadenosine, 3-methyladenosine, and N⁶-methyladenosine, 8-hydroxyguanosine, O⁶-methylguanosine, O⁶-ethylguanosine, O⁶-isopropylguanosine, 3,N²-ethenoguanosine, O⁶-alkylguanosine, 8-oxo-guanosine, 2,N³-ethenoguanosine, and 8-aminoguanosine.

11. (Canceled)

12. (Previously presented) The method of claim 1, wherein the virus is a pestivirus.

13. (Original) The method of claim 1, wherein the polymerase is a human polymerase II.
14. (Original) The method of claim 1, wherein the cell is in cell culture.
15. (Original) The method of claim 1, wherein the cell is in an animal.
16. (Original) The method of claim 1, wherein increasing the mutation rate of the virus produces a progressive loss of viability of the virus.
17. (Original) The method of claim 1, comprising administration of more than one species of RNA nucleoside analog to the virally infected cell.
18. (Currently amended) A The method of ~~claim 1~~ increasing the mutation rate of a virus, comprising administering an RNA nucleoside analog to a virally infected cell, wherein the analog is incorporated by a polymerase into an RNA copy of a genomic nucleic acid encoding the virus, said analog replacing a first natural occurring nucleotide having a first complementary nucleotide wherein said analog complements a second nucleotide which is other than the first nucleotide, thereby inducing the virus to mutate, wherein the virus is an RNA virus selected from the group consisting of hepatitis C, coronavirus, influenza, respiratory syncytial virus, BVDV, and dengue fever.
- 19-44. (Canceled)
45. (Previously presented) A method of increasing the mutation rate of a virus in an animal comprising administering to the animal a therapeutically effective dose of a mutagenic ribonucleoside analog composition wherein the analog is one that, in a cell infected with a virus of interest, is incorporated by a polymerase into an RNA copy of a genomic nucleic acid encoding the virus, said analog replacing a first natural occurring nucleotide having a first complementary nucleotide wherein said analog complements a second nucleotide which is other than the first nucleotide together with a pharmaceutically acceptable carrier.

46. (Original) The method of claim 45, wherein the nucleoside analog is selected from the group consisting of N⁴-aminocytidine, N¹-methyl-N⁴-aminocytidine, 3,N⁴-ethenocytidine, 3-methylcytidine, 5-hydroxycytidine, N⁴-dimethylcytidine, 5-(2-hydroxyethyl)cytidine, 5-chlorocytidine, 5-bromocytidine, N⁴-methyl-N⁴-aminocytidine, 5-aminocytidine, 5-nitrosocytidine, 5-(hydroxyalkyl)-cytidine, 5-(thioalkyl)-cytidine and cytidine glycol, 5-hydroxyuridine, 3-hydroxyethyluridine, 3-methyluridine, O²-methyluridine, O²-ethyluridine, 5-aminouridine, O⁴-methyluridine, O⁴-ethyluridine, O⁴-isobutyluridine, O⁴-alkyluridine, 5-nitrosouridine, 5-(hydroxyalkyl)-uridine, and 5-(thioalkyl)-uridine, 1,N⁶-ethenoadenosine, 3-methyladenosine, and N⁶-methyladenosine, 8-hydroxyguanosine, O⁶-methylguanosine, O⁶-ethylguanosine, O⁶-isopropylguanosine, 3,N²-ethenoguanosine, O⁶-alkylguanosine, 8-oxo-guanosine, 2,N³-ethenoguanosine, and 8-aminoguanosine.

47. (Original) The method of claim 45, wherein the RNA nucleoside analog is incorporated by a polymerase present in virally infected cells of the animal into an RNA copy of a genomic nucleic acid of the virus with an efficiency at least about 0.1% that of a naturally occurring complementary nucleic acid.

48. (Previously presented) The method of claim 45, wherein the animal is a human patient infected with the virus selected from the group consisting of HIV-1, HIV-2, HTLV-1, HTLV-2, hepatitis A, hepatitis B, hepatitis C, and dengue fever virus.

49. (Original) The method of claim 45, wherein the animal is a human patient having a disease selected from the group consisting of AIDS, hepatitis B, hepatitis C, T-cell leukemia.

50. (Original) The method of claim 45, the animal having a disease selected from the group consisting of feline leukemia virus, feline immunodeficiency virus, BVDV, or vesicular stomatitis virus.

51-65. (Canceled)

Appl. No. 09/522,373
Amdt. dated October 27, 2003
Reply to Office Action of June 4, 2003

PATENT

66. (Original) The method of claim 1, wherein the RNA nucleoside analog is an enantio-specific nucleoside analog.

67. (Canceled)